

# SOME IDENTITIES ON BERNSTEIN POLYNOMIALS ASSOCIATED WITH $q$ -EULER POLYNOMIALS

A. BAYAD, T. KIM, B. LEE, AND S.-H. RIM

**Abstract** In this paper we investigate some interesting properties of the  $q$ -Euler polynomials. The purpose of this paper is to give some relationships between Bernstein and  $q$ -Euler polynomials which are derived by the  $p$ -adic integral representation of the Bernstein polynomials associated with  $q$ -Euler polynomials.

## 1. INTRODUCTION

Let  $p$  be a fixed odd prime number. Throughout this paper  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$  and  $\mathbb{C}_p$  denote the ring of  $p$ -adic integers, the field of  $p$ -adic numbers and the field of  $p$ -adic completion of the algebraic closure of  $\mathbb{Q}_p$ , respectively (see [1–15]). Let  $\mathbb{N}$  be the set of natural numbers and  $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$ . The normalized  $p$ -adic absolute value is defined by  $|p|_p = \frac{1}{p}$ . As an indeterminate, we assume that  $q \in \mathbb{C}_p$  with  $|1 - q|_p < 1$ . Let  $UD(\mathbb{Z}_p)$  be the space of uniformly differentiable function on  $\mathbb{Z}_p$ . For  $f \in UD(\mathbb{Z}_p)$ , the  $p$ -adic invariant integral on  $\mathbb{Z}_p$  is defined by

$$\begin{aligned} I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) &= \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) \mu_{-1}(x + p^N \mathbb{Z}_p) \\ &= \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) (-1)^x, \quad (\text{see [7, 8, 9, 10]}). \end{aligned} \quad (1)$$

For  $n \in \mathbb{N}$ , we can derive the following integral equation from (1):

$$I_{-1}(f_n) = (-1)^n \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) + 2 \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l), \quad (2)$$

where  $f_n(x) = f(x+n)$  (see [7, 8, 9, 10, 11]). As well known definition, the Euler polynomials are given by the generating function as follows:

$$\frac{2}{e^t + 1} e^{xt} = e^{E(x)t} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}, \quad (\text{see [7, 8, 9, 10, 11, 12, 13, 5, 15, 14, 3]}), \quad (3)$$

with usual convention about replacing  $E^n(x)$  by  $E_n(x)$ . In the special case  $x = 0$ ,  $E_n(0) = E_n$  are called the  $n$ -th Euler numbers. From (3), we can derive the following recurrence formula for Euler numbers

$$E_0 = 1, (E+1)^n + E_n = \begin{cases} 2 & \text{if } n = 0, \\ 0 & \text{if } n > 0, \end{cases} \quad (\text{see [12]}).$$

---

*2000 Mathematics Subject Classification* : 11B68, 11B73, 41A30.

*Key words and phrases* : Bernstein polynomial, Euler numbers and polynomials,  $q$ -Euler numbers and polynomials, fermionic  $p$ -adic integral.

with usual convention about replacing  $E^n$  by  $E_n$ . By the definitions of Euler numbers and polynomials, we get

$$E_n(x) = (E + x)^n = \sum_{l=0}^n \binom{n}{l} x^{n-l} E_l, \quad (\text{see } [7, 8, 9, 10, 11, 12, 13, 5, 15, 14, 3]).$$

Let  $C[0, 1]$  denote the set of continuous functions on  $[0, 1]$ . For  $f \in C[0, 1]$ , Bernstein introduced the following well-known linear positive operator in the field of real numbers  $\mathbb{R}$  :

$$\mathbb{B}_n(f|x) = \sum_{k=0}^n f\left(\frac{k}{n}\right) \binom{n}{k} x^k (1-x)^{n-k} = \sum_{k=0}^n f\left(\frac{k}{n}\right) B_{k,n}(x),$$

where  $\binom{n}{k} = \frac{n(n-1)\cdots(n-k+1)}{k!} = \frac{n!}{k!(n-k)!}$  (see [1, 2, 7, 11, 12, 14]). Here,  $\mathbb{B}_n(f|x)$  is called the Bernstein operator of order  $n$  for  $f$ . For  $k, n \in \mathbb{Z}_+$ , the Bernstein polynomials of degree  $n$  are defined by

$$B_{k,n}(x) = \binom{n}{k} x^k (1-x)^{n-k}, \quad \text{for } x \in [0, 1]. \quad (4)$$

In this paper, we study the properties of  $q$ -Euler numbers and polynomials. From these properties we investigate some identities on the  $q$ -Euler numbers and polynomials. Finally, we give some relationships between Bernstein and  $q$ -Euler polynomials, which are derived by the  $p$ -adic integral representation of the Bernstein polynomials associated with  $q$ -Euler polynomials.

## 2. $q$ -EULER NUMBERS AND POLYNOMIALS

In this section we assume that  $q \in \mathbb{C}_p$  with  $|1 - q|_p < 1$ . Let  $f(x) = q^x e^{xt}$ . From (1) and (2), we have

$$\int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \frac{2}{qe^t + 1} e^{xt}. \quad (5)$$

Now, we define the  $q$ -Euler numbers as follows:

$$\frac{2}{qe^t + 1} = e^{E_q t} = \sum_{n=0}^{\infty} E_{n,q} \frac{t^n}{n!}, \quad (6)$$

with the usual convention about replacing  $E_q^n$  by  $E_{n,q}$ .

By (6), we easily get

$$E_{0,q} = \frac{2}{q+1}, \text{ and } q(E_q + 1)^n + E_{n,q} = \begin{cases} 2 & \text{if } n = 0, \\ 0 & \text{if } n > 0, \end{cases} \quad (7)$$

with usual convention about replacing  $E_q^n$  by  $E_{n,q}$ .

We note that

$$\frac{2}{qe^t + 1} = \frac{2}{e^t + q^{-1}} \cdot \frac{2}{1+q} = \frac{2}{1+q} \sum_{n=0}^{\infty} H_n(-q^{-1}) \frac{t^n}{n!}, \quad (8)$$

where  $H_n(-q^{-1})$  is the  $n$ -th Frobenius-Euler numbers.

From (5), (6) and (8), we have

$$\int_{\mathbb{Z}_p} q^x e^{xt} d\mu_{-1}(x) = E_{n,q} = \frac{2}{1+q} H_n(-q^{-1}), \text{ for } n \in \mathbb{Z}_+. \quad (9)$$

Now, we consider the  $q$ -Euler polynomials as follows:

$$\frac{2}{qe^t + 1}e^{xt} = e^{E_q(x)t} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}, \quad (10)$$

with the usual convention  $E_q^n(x)$  by  $E_{n,q}(x)$ .

From (2), (5) and (10), we get

$$\int_{\mathbb{Z}_p} q^x e^{(x+y)t} d\mu_{-1}(y) = \frac{2}{qe^t + 1}e^{xt} = \sum_{n=0}^{\infty} E_{n,q}(x) \frac{t^n}{n!}. \quad (11)$$

By comparing the coefficients on the both sides of (10) and (11), we get the following Witt's formula for the  $q$ -Euler polynomials as follows:

$$\int_{\mathbb{Z}_p} q^y (x+y)^n d\mu_{-1}(y) = E_{n,q}(x) = \sum_{l=0}^n \binom{n}{l} x^{n-l} E_{l,q}. \quad (12)$$

From (10) and (12), we can derive the following equation:

$$\frac{2q}{qe^t + 1}e^{(1-x)t} = \frac{2}{1+q^{-1}e^{-t}}e^{-xt} = \sum_{n=0}^{\infty} E_{n,q^{-1}}(x)(-1)^n \frac{t^n}{n!}. \quad (13)$$

By (10) and (13), we obtain the following reflection symmetric property for the  $q$ -Euler polynomials.

**Theorem 1.** For  $n \in \mathbb{Z}_+$ , we have

$$(-1)^n E_{n,q^{-1}}(x) = q E_{n,q}(1-x).$$

From (9), (10), (11) and (12), we can derive the following equation: for  $n \in \mathbb{N}$ ,

$$\begin{aligned} E_{n,q}(2) &= (E_q + 1 + 1)^n = \sum_{l=0}^n \binom{n}{l} E_{l,q}(1) \\ &= E_{0,q} + \frac{1}{q} \sum_{l=1}^n \binom{n}{l} q E_{l,q}(1) = \frac{2}{1+q} - \frac{1}{q} \sum_{l=1}^n \binom{n}{l} E_{l,q} \\ &= \frac{2}{1+q} + \frac{2}{q(1+q)} - \frac{1}{q} \sum_{l=0}^n \binom{n}{l} E_{l,q} \\ &= \frac{2}{q} - \frac{1}{q^2} q E_{n,q}(1) = \frac{2}{q} + \frac{1}{q^2} E_{n,q}, \text{ by using recurrence formula (7).} \end{aligned}$$

Therefore, we obtain the following theorem.

**Theorem 2.** For  $n \in \mathbb{N}$ , we have

$$q E_{n,q}(2) = 2 + \frac{1}{q} E_{n,q}.$$

By using (9) and (12), we get

$$\begin{aligned} \int_{\mathbb{Z}_p} q^{-x} (1-x)^n d\mu_{-1}(x) &= (-1)^n \int_{\mathbb{Z}_p} q^{-x} (x-1)^n d\mu_{-1}(x) \\ &= (-1)^n E_{n,q^{-1}}(-1) = q \int_{\mathbb{Z}_p} (x+2)^n d\mu_{-1}(x) = q \left( \frac{2}{q} + \frac{1}{q^2} E_{n,q} \right) \\ &= 2 + \frac{1}{q} E_{n,q} = 2 + \frac{1}{q} \int_{\mathbb{Z}_p} x^n q^x d\mu_{-1}(x), \text{ for } n > 0. \end{aligned}$$

Therefore, we obtain the following theorem.

**Theorem 3.** *For  $n \in \mathbb{N}$ , we have*

$$\int_{\mathbb{Z}_p} q^{-x} (1-x)^n d\mu_{-1}(x) = 2 + \frac{1}{q} \int_{\mathbb{Z}_p} x^n q^x d\mu_{-1}(x).$$

By using Theorem 3, we will study for the  $p$ -adic integral representation on  $\mathbb{Z}_p$  of the Bernstein polynomials associated with  $q$ -Euler polynomials in the next section.

### 3. BERNSTEIN POLYNOMIALS ASSOCIATED WITH $q$ -EULER NUMBERS AND POLYNOMIALS

Now, we take the  $p$ -adic integral on  $\mathbb{Z}_p$  for the Bernstein polynomials in (4) as follows:

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n}(x) q^x d\mu_{-1}(x) &= \int_{\mathbb{Z}_p} \binom{n}{k} x^k (1-x)^{n-k} q^x d\mu_{-1}(x) \\ &= \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^{n-k-j} \int_{\mathbb{Z}_p} x^{n-j} q^x d\mu_{-1}(x) \\ &= \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^{n-k-j} E_{n-j,q} \\ &= \binom{n}{k} \sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^j E_{k+j,q}, \quad \text{where } n, k \in \mathbb{Z}_+. \end{aligned} \tag{14}$$

By the definition of Bernstein polynomials, we see that

$$B_{k,n}(x) = B_{n-k,n}(1-x), \quad \text{where } n, k \in \mathbb{Z}_+. \tag{15}$$

Let  $n, k \in \mathbb{Z}_+$  with  $n > k$ . Then, by (15), we get

$$\begin{aligned} \int_{\mathbb{Z}_p} q^x B_{k,n}(x) d\mu_{-1}(x) &= \int_{\mathbb{Z}_p} q^x B_{n-k,n}(1-x) d\mu_{-1}(x) \\ &= \binom{n}{n-k} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} \int_{\mathbb{Z}_p} (1-x)^{n-j} q^x d\mu_{-1}(x) \\ &= \binom{n}{k} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} \left( 2 + q \int_{\mathbb{Z}_p} x^{n-j} q^x d\mu_{-1}(x) \right) \\ &= \binom{n}{k} \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} (2 + q E_{n-j,q^{-1}}) \\ &= \begin{cases} 2 + q E_{n,q^{-1}} & \text{if } k = 0, \\ \binom{n}{k} q \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} E_{n-j,q^{-1}} & \text{if } k > 0. \end{cases} \end{aligned}$$

Thus, we obtain the following theorem.

**Theorem 4.** For  $n, k \in \mathbb{Z}_+$  with  $n > k$ , we have

$$\int_{\mathbb{Z}_p} q^{1-x} B_{k,n}(x) d\mu_{-1}(x) = \begin{cases} 2q + E_{n,q} & \text{if } k = 0, \\ \sum_{j=0}^k \binom{n}{k} \binom{k}{j} (-1)^{k-j} E_{n-j,q} & \text{if } k > 0. \end{cases}$$

By (14) and Theorem 4, we get the following corollary.

**Corollary 5.** For  $n, k \in \mathbb{Z}_+$  with  $n > k$ , we have

$$\sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^j E_{k+j,q^{-1}} = \begin{cases} 2 + \frac{1}{q} E_{n,q} & \text{if } k = 0, \\ \sum_{j=0}^k \binom{k}{j} (-1)^{k-j} \frac{1}{q} E_{n-j,q} & \text{if } k > 0. \end{cases}$$

For  $m, n, k \in \mathbb{Z}_+$  with  $m + n > 2k$ . Then we get

$$\begin{aligned} & \int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) q^{-x} d\mu_{-1}(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{j=0}^{2k} \binom{2k}{j} (-1)^{j+2k} \int_{\mathbb{Z}_p} q^{-x} (1-x)^{n+m-j} d\mu_{-1}(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{j=0}^{2k} \binom{2k}{j} (-1)^{j+2k} q \int_{\mathbb{Z}_p} (x+2)^{n+m-j} q^x d\mu_{-1}(x) \\ &= \binom{n}{k} \binom{m}{k} \sum_{j=0}^{2k} \binom{2k}{j} (-1)^{j+2k} q \left( \frac{2}{q} + \frac{1}{q^2} E_{n+m-j,q} \right) \\ &= \begin{cases} 2 + \frac{1}{q} E_{n+m,q} & \text{if } k = 0, \\ \binom{n}{k} \binom{m}{k} \sum_{j=0}^{2k} \binom{2k}{j} (-1)^{j+2k} \frac{1}{q} E_{n+m-j,q} & \text{if } k > 0. \end{cases} \end{aligned}$$

Therefore, we obtain the following theorem.

**Theorem 6.** For  $m, n, k \in \mathbb{Z}_+$  with  $m + n > 2k$ , we have

$$\begin{aligned} & \int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) q^{1-x} d\mu_{-1}(x) \\ &= \begin{cases} 2q + E_{n+m,q} & \text{if } k = 0, \\ \binom{n}{k} \binom{m}{k} \sum_{j=0}^{2k} \binom{2k}{j} (-1)^{j+2k} E_{n+m-j,q} & \text{if } k > 0. \end{cases} \end{aligned}$$

By using binomial theorem, for  $m, n, k \in \mathbb{Z}_+$ , we get

$$\int_{\mathbb{Z}_p} B_{k,n}(x) B_{k,m}(x) q^{1-x} d\mu_{-1}(x) \quad (16)$$

$$= \binom{n}{k} \binom{m}{k} \sum_{j=0}^{n+m-2k} \binom{n+m-2k}{j} (-1)^j \int_{\mathbb{Z}_p} x^{j+2k} q^{1-x} d\mu_{-1}(x) \quad (17)$$

$$= q \binom{n}{k} \binom{m}{k} \sum_{j=0}^{n+m-2k} \binom{n+m-2k}{j} (-1)^j E_{j+2k, q^{-1}}. \quad (18)$$

By comparing the coefficients on the both sides of (16) and Theorem 6, we obtain the following corollary.

**Corollary 7.** *Let  $m, n, k \in \mathbb{Z}_+$  with  $m + n > 2k$ . Then we get*

$$\begin{aligned} & \sum_{j=0}^{n+m-2k} \binom{n+m-2k}{j} (-1)^j E_{j+2k, q^{-1}} \\ &= \begin{cases} 2 + \frac{1}{q} E_{n+m, q} & \text{if } k = 0, \\ \frac{1}{q} \sum_{j=0}^{2k} \binom{2k}{j} (-1)^{j+2k} E_{n+m-j, q} & \text{if } k > 0. \end{cases} \end{aligned}$$

For  $s \in \mathbb{N}$ , let  $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$  with  $n_1 + n_2 + \dots + n_s > sk$ . By induction, we get

$$\begin{aligned} \int_{\mathbb{Z}_p} B_{k,n_1}(x) \cdots B_{k,n_s}(x) q^{-x} d\mu_{-1}(x) &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \int_{\mathbb{Z}_p} x^{sk} (1-x)^{n_1+\dots+n_s} q^{-x} d\mu_{-1}(x) \\ &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{j=0}^{sk} \binom{sk}{j} (-1)^{sk+j} \int_{\mathbb{Z}_p} (1-x)^{n_1+\dots+n_s-j} q^{-x} d\mu_{-1}(x) \\ &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{j=0}^{sk} \binom{sk}{j} (-1)^{sk+j} q \int_{\mathbb{Z}_p} (x+2)^{n_1+\dots+n_s-j} q^x d\mu_{-1}(x) \\ &= \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{j=0}^{sk} \binom{sk}{j} (-1)^{sk+j} q \left( \frac{2}{q} + \frac{1}{q^2} E_{n_1+\dots+n_s-j, q} \right) \\ &= \begin{cases} 2 + \frac{1}{q} E_{n_1+\dots+n_s, q} & \text{if } k = 0, \\ \left( \prod_{i=1}^s \binom{n_i}{k} \right) \frac{1}{q} \sum_{j=0}^{sk} \binom{sk}{j} (-1)^{sk+j} E_{n_1+\dots+n_s-j, q} & \text{if } k > 0. \end{cases} \end{aligned}$$

Therefore we obtain the following theorem.

**Theorem 8.** Let  $s \in \mathbb{N}$ . For  $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$  with  $n_1 + n_2 + \dots + n_s > sk$ , we have

$$\begin{aligned} & \int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k, n_i}(x) \right) q^{1-x} d\mu_{-1}(x) \\ &= \begin{cases} 2q + E_{n_1+n_2+\dots+n_s, q} & \text{if } k = 0, \\ \left( \prod_{i=1}^s \binom{n_i}{k} \right) \sum_{j=0}^{sk} \binom{sk}{j} (-1)^{sk+j} E_{n_1+n_2+\dots+n_s-j, q} & \text{if } k > 0. \end{cases} \end{aligned}$$

For  $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$  by binomial theorem, we get

$$\begin{aligned} & \int_{\mathbb{Z}_p} \left( \prod_{i=1}^s B_{k, n_i}(x) \right) q^{-x} d\mu_{-1}(x) \tag{19} \\ &= \binom{n_1}{k} \dots \binom{n_s}{k} \sum_{j=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{j} (-1)^j \int_{\mathbb{Z}_p} x^{j+sk} q^{-x} d\mu_{-1}(x) \\ &= \binom{n_1}{k} \dots \binom{n_s}{k} \sum_{j=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{j} (-1)^j E_{j+sk, q^{-1}}. \end{aligned}$$

By using (19) and Theorem 8, we obtain the following corollary.

**Corollary 9.** Let  $s \in \mathbb{N}$ . For  $n_1, n_2, \dots, n_s, k \in \mathbb{Z}_+$  with  $n_1 + n_2 + \dots + n_s > sk$ , we have

$$\begin{aligned} & \sum_{j=0}^{n_1+\dots+n_s-sk} \binom{n_1+\dots+n_s-sk}{j} (-1)^j E_{j+sk, q^{-1}} \\ &= \begin{cases} 2 + \frac{1}{q} E_{n_1+n_2+\dots+n_s, q} & \text{if } k = 0, \\ \frac{1}{q} \sum_{j=0}^{sk} \binom{sk}{j} (-1)^{sk+j} E_{n_1+n_2+\dots+n_s-j, q} & \text{if } k > 0. \end{cases} \end{aligned}$$

## REFERENCES

- [1] M. Acikgoz, S. Araci, *A study on the integral of the product of several type Bernstein polynomials*, IST Transaction of Applied Mathematics-Modelling and Simulation, 2010, Vol. 1, No.1(2), ISBN 1913-8342, 10–14.
- [2] S. Bernstein, *Démonstration du théorème de Weierstrass, fondée sur le calcul des probabilités*, Commun. Soc. Math. Kharkow **13** (1912), 1–2.
- [3] I. N. Cangul, V. Kurt, H. Ozden, Y. Simsek, *On the higher-order  $w$ - $q$ -Genocchi numbers*, Adv. Stud. Contemp. Math. **19** (2009), 39–57.
- [4] N. K. Govil, V. Gupta, *Convergence of  $q$ -Meyer-König-Zeller-Durrmeyer operators*, Adv. Stud. Contemp. Math. **19** (2009), 97–108.
- [5] L.C. Jang, W-J. Kim, Y. Simsek, *A study on the  $p$ -adic integral representation on  $\mathbb{Z}_p$  associated with Bernstein and Bernoulli polynomials*, Advances in difference Equations, Vol 2010 (2010), Article ID 163219, 6pp.
- [6] K. I Joy, *Bernstein polynomials*, On-line Geometric Modelling Notes, [http://en.wikipedia.org/wiki/Bernstein\\_polynomials](http://en.wikipedia.org/wiki/Bernstein_polynomials).

- [7] T. Kim, *Barnes-type multiple  $q$ -zeta functions and  $q$ -Euler polynomials*, J. Physics A : Math. Theor., **43** (2010), 255201, 11pp.
- [8] T. Kim, J. Choi, Y. H.Kim, L. Jang, *On  $p$ -adic analogue of  $q$ -Bernstein polynomials and related integrals*, Discrete Dynamics in Nature and Society, Vol 2010 (2010), Article ID 17930, 3 pp.
- [9] T. Kim, J. Choi, Y. H.Kim,  *$q$ -Bernstein polynomials associated with  $q$ -Stirling numbers and Carlit's  $q$ -Bernoulli numbers*, Abstract and Applied Analysis (in press).
- [10] T. Kim, *Some identities on the  $q$ -Euler polynomials of higher order order and  $q$ -Stirling numbers by fermionic  $p$ -adic integral on  $\mathbb{Z}_p$* , Russian Journal of Mathematical physics 16 (2009), 484–491.
- [11] T. Kim, *Note on the Euler numbers and polynomials*, Adv. Stud. Contemp. Math. **19** (2008), 131–136.
- [12] T. Kim, B. Lee, *Some identities of the Frobenius-Euler polynomials*, Abstract and Applied Analysis 2009 (2009), Article ID 639439, 7 pp.
- [13] T. Kim, *Symmetry of power sum polynomials and multivariate fermionic  $p$ -adic invariant integral on  $\mathbb{Z}_p$* , Russian Journal of Mathematical physics 16 (2009), 484–491.
- [14] V. Kurt, *A further symmetric relation on the analogue of the Apostol-Bernoulli and the analogue of the Apostol-Genocchi polynomials*, Appl. Math. Sci. **3** (2009), 53–56.
- [15] S.-H. Rim, J.-H. Jin, E.-J. Moon, S.-J. Lee, *On multiple interpolation functions of the  $q$ -Genocchi polynomials*, Journal of inequalities and Applications, Vol 2010 (2010), Article ID 351419, 13 pp..

ABDELMEJID BAYAD. DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ D'EVRY VAL D'ESSONNE, BD. F. MITTERRAND, 91025 EVRY CEDEX, FRANCE,  
*E-mail address:* abayad@maths.univ-evry.fr

TAEKYUN KIM. DIVISION OF GENERAL EDUCATION-MATHEMATICS, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA,  
*E-mail address:* tkkim@kw.ac.kr

DEPARTMENT OF WIRELESS COMMUNICATIONS ENGINEERING,, KWANGWOON UNIVERSITY, SEOUL 139-701, REPUBLIC OF KOREA,  
*E-mail address:* bjlee@kw.ac.kr

DEPARTMENT OF MATHEMATICS EDUCATION,, KYUNGPOOK NATIONAL UNIVERSITY, TAEGU 702-701, REPUBLIC OF KOREA,  
*E-mail address:* shrim@knu.ac.kr